



# Sulfoximines for Drug Design

## Introduction


Incorporation of a sulfoximine group into bioactive compounds often improves their ADME/Tox profile, and enhances potency. Moreover, the moiety of sulfoximine is chemically and metabolically stable. NH-sulfoximines can serve as the both hydrogen bond donors and acceptors at the same time.<sup>1-6</sup> Production and commercialization of the building blocks that already contain a sulfoximine group allow significant accelerating discovery of drug candidates. Herein we have designed and synthesized a library of sulfoximines for drug design.



## Design



**We offer** >100 unique sulfoximines on a 5-50 g scale from stock.



## References

- J. A. Sirvent et al. *ChemMedChem*, 2017, 487.
- M. Frings et al. *Eur. J. Med. Chem.* 2017, 225.
- U. Lücking. *Angew. Chem. Int. Ed.* 2013, 9399.
- S. Park et al. *Bioorg. Med. Chem. Lett.* 2011, 4888.
- J. A. Bull et al. *Synlett* 2017, 28, 2525.
- D. P. Walker et al. *Bioorg. Med. Chem. Lett.* 2009, 3253.



ご興味があるものがあれば、こちらまでお問い合わせ下さい!!



E-mail : info@namiki-s.co.jp  
TEL(本社) : 03-3354-4026  
TEL(大阪支店) : 06-6231-5444